

SPECIMEN

Status®

Herbicide

For use in field corn grown for grain, seed, or silage; and popcorn

A broad-spectrum corn herbicide, in a safened formulation, specifically designed for use alone or with glyphosate combinations

Active Ingredients:

sodium salt of diflufenzopyr: 2-(1-[[[3,5-difluorophenylamino]carbonyl]-hydrazono]ethyl)-3-pyridinecarboxylic acid, sodium salt*	17.1%
sodium salt of dicamba: 3,6-dichloro-2-methoxybenzoic acid, sodium salt**	44.0%

Other Ingredients: 38.9%

Total: 100.0%

* This product contains 16% 2-(1-[[[3,5-difluorophenylamino]carbonyl]-hydrazono]ethyl)-3-pyridinecarboxylic acid (diflufenzopyr).

** This product contains 40% 3,6-dichloro-2-methoxybenzoic acid.

EPA Reg. No. 7969-242

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See inside for complete **First Aid, Precautionary Statements, Directions For Use, Conditions of Sale and Warranty**, and state-specific crop and/or use site restrictions.

In case of an emergency endangering life or property involving this product, call day or night 1-800-832-HELP (4357).

Net Contents:

FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes; then continue rinsing. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to do so by a poison control center or doctor. • DO NOT give anything to an unconscious person.
If on skin	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact BASF Corporation for emergency medical treatment information: 1-800-832-HELP (4357).</p>	

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Harmful if swallowed or absorbed through skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Personal Protective Equipment (PPE)

Some materials that are chemically resistant to this product are natural rubber and nitrile rubber.

All mixers, loaders, applicators, and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear (goggles, face shield, or safety glasses)

See **Engineering Controls** for additional requirements and exceptions.

Follow the manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Pilots must use cockpits in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)].

Environmental Hazards

DO NOT apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate. This chemical is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Groundwater and Surface Water Protection

Point-source contamination. To prevent point-source contamination, **DO NOT** mix/load this pesticide product within 50 feet of wells (including abandoned wells and drainage wells), sinkholes, perennial or intermittent streams and rivers, and natural or impounded lakes and reservoirs. **DO NOT** apply pesticide product within 50 feet of wells. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pad or properly diked mixing/loading areas as described below.

Mixing, loading, rinsing, or washing operations performed within 50 feet of a well are allowed only when conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be on or move across the pad. The pad must be self-contained to prevent surface water flow over or from the pad. The pad capacity must be maintained at 110% that of the largest pesticide container or application equipment used on the pad and have

sufficient capacity to contain all product spills, equipment or container leaks, equipment washwaters, and rainwater that may fall on the pad. The containment capacity does not apply to vehicles delivering pesticide shipments to the mixing/loading site. States may have in effect additional requirements regarding wellhead setbacks and operational containment.

Care must be taken when using this product to prevent:

- Back-siphoning into wells
- Spills
- Improper disposal of excess pesticide, spray mixtures, or rinsates.

Check valves or antisiphoning devices must be used on all mixing equipment.

Movement by surface runoff or through soil

- **DO NOT** apply under conditions which favor runoff.
- **DO NOT** apply to impervious substrates such as paved or highly compacted surfaces in areas with high potential for groundwater contamination. Groundwater contamination may occur in areas where soils are permeable or coarse and groundwater is near the surface.
- **DO NOT** apply to soils classified as sand with less than 3% organic matter and where groundwater depth is shallow. To minimize the possibility of groundwater contamination, carefully follow the application rate.

Movement by water erosion of treated soil

- **DO NOT** apply or incorporate this product through any type of irrigation equipment or by flood or furrow irrigation. Ensure treated areas have received at least 1/2-inch rainfall (or irrigation) before using tailwater for subsequent irrigation of other fields.

Endangered Species

The use of any pesticide in a manner that may kill or otherwise harm an endangered species or adversely modify their habitat is a violation of federal law.

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

All applicable directions, restrictions, precautions, and **Conditions of Sale and Warranty** are to be followed.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS), 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the WPS.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **24 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls worn over short-sleeved shirt and short pants
- Waterproof gloves
- Chemical-resistant footwear plus socks
- Chemical-resistant headgear for overhead exposure
- Protective eyewear (goggles, face shield, or safety glasses)

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store product in original container only in a cool, dry place. **DO NOT** store this product under wet conditions. Avoid cross-contamination with other pesticides.

Pesticide Disposal

Wastes resulting from the use of this product must be disposed of on-site or at an approved waste disposal facility. Improper disposal of excess pesticide, spray mix, or rinsate is a violation of federal law. If these wastes cannot be disposed of according to label instructions, contact the state agency responsible for pesticide regulation or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Handling

Nonrefillable Container. DO NOT reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Triple rinse containers small enough to shake (capacity ≤ 50 pounds) as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Triple rinse containers too large to shake (capacity > 50 pounds) as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or mix tank. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

In Case of Emergency

In case of large-scale spillage regarding this product, call:

- CHEMTREC 1-800-424-9300
- BASF Corporation 1-800-832-HELP (4357)

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment
- Your local poison control center (hospital)
- BASF Corporation 1-800-832-HELP (4357)

Steps to take if this material is released into the environment or spilled:

- Wear Personal Protective Equipment (PPE) and avoid exposure when managing a spill. (See **Precautionary Statements** section of this label for required PPE.)
- Spillage or leakage should be contained, carefully swept up, and collected for disposal. Wash area of spill with detergent.
- Remove contaminated clothing, and wash affected skin areas with soap and water. Wash clothing before reuse.

Product Information

Status® herbicide is a selective postemergence herbicide for control of annual broadleaf weeds, control or suppression of many perennial broadleaf weeds, and suppression of annual grass weeds found in field corn grown for grain, seed, or silage, and popcorn. Use **Status** sequentially or tank mix with a grass herbicide for a complete weed control program (refer to **Tank Mixing Information**).

Weeds Controlled

Status will provide postemergence control of annual and biennial broadleaf weeds and control or suppression of many perennial broadleaf weeds including ALS-resistant¹ and triazine-resistant biotypes.

Status provides suppression of annual grass weeds at appropriate rates. Emerged grass up to 3-inches tall will cease growing but may remain green for weeks after application. Regrowth of grass is limited when crop canopies over row middles.

¹ALS (acetolactate synthase)-resistant weeds include those weeds resistant to the sulfonylurea, imidazolinone, and/or sulfonamide family of herbicides.

Table 1. Weeds Controlled or Suppressed

Common Name	Scientific Name
Annual Broadleaf Weeds	
Amaranth, Palmer	<i>Amaranthus palmeri</i>
Beggarweed, Florida	<i>Desmodium tortuosum</i>
Buckwheat, wild	<i>Polygonum convolvulus</i>
Buffalobur	<i>Solanum rostratum</i>
Burcucumber	<i>Sicyos angulatus</i>
Carpetweed	<i>Mollugo verticillata</i>
Chickweed, common	<i>Stellaria media</i>

(continued)

Table 1. Weeds Controlled or Suppressed (continued)

Common Name	Scientific Name
Annual Broadleaf Weeds (continued)	
Cocklebur, common	<i>Xanthium strumarium</i>
Croton, tropic	<i>Croton glandulosus</i>
Devil's claw	<i>Proboscidea louisianica</i>
Henbit	<i>Lamium amplexicaule</i>
Jimsonweed	<i>Datura stramonium</i>
Knotweed, prostrate	<i>Polygonum aviculare</i>
Kochia	<i>Kochia scoparia</i>
Ladysthumb	<i>Polygonum persicaria</i>
Lambsquarters, common	<i>Chenopodium album</i>
Lettuce, prickly	<i>Lactuca serriola</i>
Mallow, common	<i>Malva neglecta</i>
Mallow, Venice	<i>Hibiscus trionum</i>
Marestail (Horseweed)	<i>Conyza canadensis</i>
Morningglory, entireleaf	<i>Ipomoea hederacea</i> var. <i>integriuscula</i>
Morningglory, ivyleaf	<i>Ipomoea hederacea</i> var. <i>hederacea</i>
Morningglory, pitted (small white)	<i>Ipomoea lacunosa</i>
Morningglory, smallflower	<i>Jacquemontia tamnifolia</i>
Morningglory, tall (common)	<i>Ipomoea purpurea</i>
Nightshade, black	<i>Solanum nigrum</i>
Nightshade, Eastern black	<i>Solanum ptycanthum</i>
Nightshade, hairy	<i>Solanum sarrachoides</i>
Pennycress, field	<i>Thlaspi arvense</i>
Pigweed, prostrate	<i>Amaranthus blitoides</i>
Pigweed, redroot	<i>Amaranthus retroflexus</i>
Pigweed, smooth	<i>Amaranthus hybridus</i>
Pigweed, spiny	<i>Amaranthus spinosus</i>
Pigweed, tumble	<i>Amaranthus albus</i>
Purslane, common	<i>Portulaca oleracea</i>
Radish, wild	<i>Raphanus raphanistrum</i>
Ragweed, common	<i>Ambrosia artemisiifolia</i>
Ragweed, giant	<i>Ambrosia trifida</i>
Sesbania, hemp	<i>Sesbania exaltata</i>
Shepherd's purse	<i>Capsella bursa-pastoris</i>
Sicklepod	<i>Cassia obtusifolia</i>
Sida, prickly (Teaweed)	<i>Sida spinosa</i>
Smartweed, Pennsylvania	<i>Polygonum pennsylvanicum</i>
Smellmelon	<i>Cucumis melo</i>
Sowthistle, annual	<i>Sonchus oleraceus</i>
Spurge, prostrate	<i>Chamaesyce humistrata</i>
Sunflower, volunteer	<i>Helianthus annuus</i>
Sunflower, wild (common)	<i>Helianthus annuus</i>
Thistle, Russian	<i>Salsola iberica</i>
Velvetleaf	<i>Abutilon theophrasti</i>
Waterhemp, common	<i>Amaranthus rudis</i>
Waterhemp, tall	<i>Amaranthus tuberculatus</i>

Table 1. Weeds Controlled or Suppressed (continued)

Common Name	Scientific Name
Perennial Broadleaf Weeds¹	
Alfalfa	<i>Medicago sativa</i>
Bindweed, field	<i>Convolvulus arvensis</i>
Bindweed, hedge	<i>Calystegia sepium</i>
Clover, white	<i>Trifolium repens</i>
Dandelion, common	<i>Taraxacum officinale</i>
Dock, broadleaf	<i>Rumex obtusifolius</i>
Dock, curly	<i>Rumex crispus</i>
Dogbane, hemp	<i>Apocynum cannabinum</i>
Horsenettle, Carolina	<i>Solanum carolinense</i>
Knapweed, spotted	<i>Centaurea maculosa</i>
Milkweed, common	<i>Asclepias syriaca</i>
Milkweed, honeyvine	<i>Ampelamus albidus</i>
Nightshade, silverleaf	<i>Solanum elaeagnifolium</i>
Plantain, broadleaf	<i>Plantago major</i>
Pokeweed	<i>Phytolacca americana</i>
Potato, volunteer	<i>Solanum tuberosum</i>
Smartweed, swamp	<i>Polygonum coccineum</i>
Sowthistle, perennial	<i>Sonchus arvensis</i>
Thistle, Canada	<i>Cirsium arvense</i>
Annual Grass Weeds²	
Barnyardgrass	<i>Echinochloa crus-galli</i>
Foxtail, giant	<i>Setaria faberi</i>
Foxtail, green	<i>Setaria viridis</i>
Foxtail, yellow	<i>Setaria glauca</i>
Johnsongrass, seedling	<i>Sorghum halepense</i>
Panicum, fall	<i>Panicum dichotomiflorum</i>
Shattercane	<i>Sorghum bicolor</i>
Signalgrass, broadleaf	<i>Urochloa platyphylla</i>

¹Partially controlled or suppressed

²**Status® herbicide** provides suppression of annual grass weeds at appropriate rates (5 ozs per acre or greater). Emerged grass weeds up to 3 inches tall will cease growing but may remain green for weeks after application. Regrowth of grass weeds is limited when corn canopies over row middles.

Mode of Action

Status is absorbed by leaves, roots, and shoots and is translocated to the growing points of sensitive weeds to provide postemergence control of emerged weeds and moderate residual control of germinating weeds. **Status** controls weeds by auxin-transport inhibition and auxin-agonist modes of action.

Treated weeds will stop growing soon after application. Broadleaf weeds will display epinastic twisting and crinkling symptoms before becoming necrotic. Suppressed grass weeds may display some epinasty and remain stunted and green.

Crop Tolerance

Corn is very tolerant to an application of **Status**. Corn growing under stress conditions, such as drought, poor fertility, or foliar damage because of hail, wind, or insects,

can show various injury symptoms that may be more pronounced if **Status® herbicide** is applied. Injury can be avoided by agronomic practices that promote good crop growth and minimize stress conditions, especially combinations of stress factors.

Coverage

Weeds must be thoroughly covered with spray. Dense leaf canopies shelter smaller weeds and can prevent adequate spray coverage.

Cultivation

Avoid disturbing (e.g. tillage or cultivating) treated areas for at least 7 days following application to allow best herbicide uptake, translocation, and weed control.

Cleaning Spray Equipment

Clean application equipment thoroughly by using a strong detergent or commercial sprayer cleaner according to the manufacturer's directions. Triple rinse the equipment before and after applying this product.

Application Instructions

Best product performance is obtained when **Status** is applied to actively growing weeds. Treated weeds will stop growing soon after application. Broadleaf weeds will display epinastic twisting and crinkling symptoms before becoming necrotic. Suppressed grass weeds may display some epinasty and remain stunted and green.

Ground Application Methods and Equipment

Status, a wettable granule formulation, can be applied using water as the spray carrier.

Water Volume. Select an appropriate spray volume that ensures adequate coverage of the target weed species. Use higher water volumes when treating dense or tall vegetation. **DO NOT** apply less than 3 gallons of spray volume per acre.

Application Equipment. Use application equipment that will provide good spray coverage of weed foliage. Exercise preventive measures to avoid drift onto nontarget areas.

Spray Drift Management

Avoiding spray drift at the application site is the responsibility of the applicator. **DO NOT** spray when conditions favor drift beyond area intended for application. Conditions which may contribute to drift include thermal inversion, wind speed and direction, spray nozzle/pressure combinations, spray droplet size, temperature/humidity, etc. Agriculturally approved drift-reducing additives are recommended. Contact your state extension agent for spray drift prevention guidelines in your area. Application equipment must be properly maintained and calibrated using appropriate carriers.

Spray Additives

Adjuvants must be used with **Status** for consistent weed control.

For best results under most conditions, combine an adjuvant with a nitrogen source.

Rainfast Period - **Status** is rainfast **4 hours** after applications when used with recommended adjuvants.

Adjuvant

- Crop oil concentrate (COC)
1 to 2 pints/A
- Methylated seed oil (MSO)
1 to 2 pints/A
- Nonionic surfactant (NIS)
0.25% volume/volume (v/v) **or** 1 quart/100 gallons

Nitrogen Source

- Ammonium sulfate (AMS)
(21% nitrogen)

5 pounds to 17 pounds/100 gallons spray mix

Before adding AMS, dissolve **Status** in the spray tank. Use high-quality AMS (spray grade) to avoid plugging nozzles.

- Urea ammonium nitrate (UAN)
(28% to 34% nitrogen)

Use at least 5 quarts/100 gallons
(1.25% v/v)

Because most nitrogen solutions are mildly corrosive to galvanized steel, mild steel, and brass spray equipment, rinse the entire spray system with water soon after use.

Compatibility Test for Tank Mix Components

1. Before mixing components, always perform a compatibility jar test. Begin with a quart-sized jar. Add components in the same order as listed in **Tank Mixing Order** section. Start with 3.5 cups of water from the intended source at the source temperature. For each dry product, add 2 teaspoons per pound of product per acre. For each liquid product, add 1 teaspoon per pint of product per acre.
2. Always cap the jar and invert 10 cycles between component additions.
3. When the components have all been added to the jar, let the solution stand for 15 minutes.
4. **Evaluate** the solution for uniformity and stability. The spray solution should not have free oil on the surface; fine particles that precipitate to the bottom; or thick (clabbered) texture. **DO NOT** use any spray solution that could clog spray nozzles.

Tank Mixing Order

1. **Water** - Begin by agitating a thoroughly clean sprayer tank 1/2 full of clean water.*
2. **Products in PVA bags** - Place any product contained in water-soluble PVA bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
3. **Water-dispersible products** (dry flowables, wetttable powders, suspension concentrates, or suspo-emulsions)
4. **Water-soluble products** (such as **Status® herbicide**) - **Status** must be fully dissolved before adding other components.
5. **Emulsifiable concentrates** (including COC and MSO).
6. **Water-soluble additives** (AMS, NIS, or UAN when applicable)
7. **Remaining quantity of water**

Maintain constant agitation during application. For more information, refer to **Tank Mixing Information**. Refer to the drift-reducing additive label for proper addition rate and mixing order.

* User may fill the spray tank from a nurse tank containing an AMS product dissolved in water. For this method, thoroughly dissolve the AMS product before adding **Status**. **Status** must be thoroughly dissolved before adding additional products or additives. Verify that the AMS pre-mix water alternative is compatible with other tank mix components.

Tank Mixing Information

Use **Status** sequentially or tank mix with other herbicides as part of a complete weed control program. Tank mix recommendations are for use only in states where the sequential or tank mix product and application site are registered. Refer to **Crop-specific Information** for more details and for specific tank mix restrictions. Local agricultural authorities may be a source of information when using other than BASF-recommended tank mixes. **It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.**

Physical incompatibility, reduced weed control, or crop injury may result from mixing **Status** with other pesticides (fungicides, herbicides, insecticides, or miticides), additives, or fertilizers.

Restrictions

- **Maximum seasonal use rate** - **DO NOT** apply more than a total of 12.5 ozs of **Status** (0.438 pound ae) per acre per season.
- **Restricted-entry Interval (REI) - 24 hours.**
- **DO NOT** apply to corn showing injury (leaf phytotoxicity or plant stunting) produced by any other prior herbicide

application because this injury may be enhanced or prolonged.

- **DO NOT** apply if corn is more than 36 inches tall, or V8 stage, or within 15 days before tassel emergence, whichever comes first.
- For sequential applications, **DO NOT** apply less than 15 days apart.
- **Preharvest interval (PHI)**
 - **DO NOT** apply within 32 days before corn forage harvest.
 - **DO NOT** apply within 72 days before popcorn, corn grain and stover harvest.
- See **Table 2. Crop-specific Restrictions**
- **DO NOT** apply through any type of irrigation system.
- **DO NOT** treat irrigation ditches or water used for crop irrigation or domestic uses.
- **DO NOT** use sprayable fluid fertilizer as the carrier for application of **Status** made after corn emergence.
- **Crop Rotation Restrictions** - **DO NOT** plant any crops within 120 days after the last application of **Status**, with the following exceptions:
 - If at least 1 inch of rainfall or overhead irrigation is received following the last application of **Status** (less than or equal to 5 ozs per acre only), alfalfa, cereal grain crops, cotton, grain sorghum, and soybeans may be planted 30 days after the rainfall/irrigation event in all states except California.
 - In the event of crop failure, corn can be replanted 7 or more days after application.

Table 2. Crop-specific Restrictions

Crop	Minimum Time from Application to Harvest (PHI) (days)	Maximum Rate per acre per Application (ozs)	Maximum Rate per acre per Year (ozs)
Corn forage	32	10.0	12.5
Corn grain or popcorn or stover	72		

Corn Use Directions

Field Corn (grown for grain, seed, or silage) and Popcorn

Before applying **Status**[®] herbicide to popcorn or seed corn, verify the selectivity of **Status** on the inbred line or hybrid with your local seed corn company to help avoid potential injury of sensitive lines.

Application Rates and Timings

Status can be applied at the timing, rates, and growth stages listed in **Table 3** in all tillage systems (e.g. conventional, no-tillage, reduced tillage).

• Postemergence Application

Status can be applied from 4-inch tall (V2) to 36-inch tall (V8) corn at rates from 5 to 10 ozs/A.

- Apply early postemergence for best weed control and crop yield potential.

• **Status** may be tank mixed with **Lightning**[®] herbicide, **Liberty**[®] herbicide, **Roundup**[®] herbicide, or glyphosate.

- Apply 2.5 to 10 ozs of **Status** with the tank mix herbicides (e.g. **Lightning**, **Liberty**, **Roundup**, or glyphosate).

- Use the minimum **Status** rate of 5 ozs/A for weeds resistant to the tank mix herbicide (ALS-resistant, glyphosate-resistant, or PPO-resistant weeds), for perennial weeds listed on **Table 1**, weeds taller than 6 inches, or weeds not controlled by the tank mix partner.

- **Lightning**, **Liberty**, **Roundup**, and glyphosate can only be used on specifically designated corn varieties (e.g. **Lightning** on **Clearfield**[®] corn, **Liberty** on **LibertyLink**[®] corn, and **Roundup** or glyphosate on **Roundup Ready**[®] corn). **DO NOT** use on corn varieties that are not labeled for use.

• Split Application

Split applications of **Status** may be made during a growing season. **DO NOT** exceed a total of 12.5 ozs of **Status** per treated acre per crop year. **DO NOT** apply less than 15 days apart.

Field Corn Tank Mixes and Sequential Uses

In addition to control of many broadleaf weed species, **Status** offers herbistatic suppression of several annual grass weeds that may enhance overall control of your grass herbicide program.

For commercial control of grass weeds, use **Status** as a sequential postemergence treatment following a preemergence grass herbicide (e.g. **G-Max Lite**[™] herbicide, **Guardsman Max**[®] herbicide, or **Outlook**[®] herbicide) or in tank mix combination with a postemergence grass herbicide (e.g. **Lightning**, **Liberty**, **Option**[®] herbicide, **Roundup**, or glyphosate).

Tank Mix Recommendations

Status may be applied sequentially or in tank mixes with other herbicides registered for use in corn with the following limitations:

- Postemergence applications of **Status** are not recommended for use in tank mixes with plant growth regulating herbicides such as products containing dicamba, 2,4-D, or clopyralid herbicides. Additionally, sequential treatments with these products should be separated by at least 15 days.
- Tank mixes with emulsifiable concentrate (EC) formulations of chloroacetamide grass herbicides (e.g. **Outlook**, **Dual II Magnum**[®] herbicide, **Harness**[®] herbicide, or **Surpass**[®] herbicide) are not recommended.

Status may be used sequentially with all soil-applied insecticides or used sequentially or in tank mixes with foliar-applied insecticides with the following limitations:

Status is not recommended for use in foliar-applied tank mixes with **Lorsban**[®] insecticide. However, sequential treatments with these products may be used if applications are separated by at least 7 days.

Table 3. Application Rates and Timing in Field Corn and Popcorn

Application Timing	Corn Stage (inches)	Status Rate ¹ Solo Application ² (ozs/A)	Status Rate ¹ Tank Mix Application ³ (ozs/A)
Postemergence	4 to 36 inches (V2 to V8)	5 to 10	2.5 to 10

¹ Increase rate of **Status** if target weeds are:

- Resistant to tank mix partner
- Perennial weeds
- Taller than 6 inches
- Weeds not controlled by tank mix partner

² Solo rate also includes tank mixes with grass herbicides with little broadleaf activity.

³ For tank mixes with **Lightning**, **Liberty**, **Roundup**, or glyphosate, refer to their respective labels for use rates.

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

Status® herbicide is specially formulated for use on crops as specified in this label. Bayer CropScience is the owner of United States patent rights to methods and compositions covering certain uses, particularly the following United States Patents: 6,486,096; 6,569,805; 5,922,646; and 5,516,750.

Clearfield, Guardsman Max, Lightning, Outlook, and Status are registered trademarks of BASF.

G-Max Lite is a trademark of BASF.

Dual II Magnum is a registered trademark of a Syngenta Group Company.

Harness, Roundup, and Roundup Ready are registered trademarks of Monsanto Technology LLC.

Liberty, LibertyLink, and Option are registered trademarks of Bayer.

Lorsban and Surpass are registered trademarks of Dow AgroSciences LLC.

© 2017 BASF Corporation
All rights reserved.

007969-00242.20170831.**NVA 2017-04-279-0017**

Based on: NVA 2015-04-279-0156
Supersedes: NVA 2009-04-279-0118

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709



We create chemistry